



This is an official
CDC Health Advisory

Distributed via Health Alert Network
January 28, 2013, 3:45PM
10276-CHA-01-28-2013-INH

Recommendations for Drug Allocation, Tuberculosis Prevention, and Patient Care During Isoniazid Shortages

Summary

Shortages of isoniazid (INH), a cornerstone drug for treating tuberculosis disease (TB) and latent Mycobacterium tuberculosis infection (LTBI), are continuing. This notice gives an update on the shortages and expands general guidance to public health officials and clinicians about how to adjust practices in response to the shortages from that outlined in CDC's December 21, 2012, Morbidity and Mortality Weekly Report (MMWR) (<http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6150a4.htm>). It also outlines national plans for restoring INH supplies and lists published guidance that could assist in making treatment decisions when INH is unavailable.

Background

U.S. INH production has been interrupted, and stocks are dwindling. Consequent to the critical shortage of 300 mg tablets that was recently reported (1), the stocks of 100 mg tablets in some localities are reportedly being depleted. An initial forecast date for restoring INH production has been extended from late January 2013 to as late as March 2013 (1,2); however, the forecasts are unstable and vary by company. One of three pharmaceutical companies supplying INH, VersaPharm Incorporated, Marietta, Georgia, has notified public health officials that it is cancelling all INH backorders and not accepting new orders.

Clinical and Public Health Recommendations

While INH is in shortage, clinicians should coordinate TB and LTBI treatment plans with their jurisdictional public health officials, who should set priorities for allocating INH stock locally. These priorities should be set according to the immediate medical needs of individual patients and a general TB control strategy:

- 1) treating patients who have TB disease,
- 2) treating LTBI patients who are diagnosed during contact tracing of contagious TB (i.e., contact investigations), and
- 3) treating LTBI patients who face the greatest likelihood of TB disease (e.g., HIV-infected patients) or the hazard of severe illness (e.g., children <5 years old).

Treatment for LTBI can be postponed when the likelihood or hazard of TB disease is low, but systems should be developed for recalling patients when INH becomes available.

The World Health Organization (WHO) recommends that "drugs for the full course of treatment are reserved for the patient at the outset of treatment" (3). However, the abrupt onset of the INH shortage means that treatment for some TB patients could be disrupted in mid-course. Decisions for continuing

treatment in these situations should be made on a case-by-case basis in consultation with a TB expert, as arranged by jurisdictional public health officials.

Fixed dose combination formulations of INH and rifampin or INH, rifampin, and pyrazinamide can be used for treating TB disease if the dosages of all the drugs are correct for the patient (4). However, at least one fixed dose combination is already on backorder, and stocks are believed to be depleted. A full treatment course of drugs should be secured before starting treatment with one of these combinations.

Regimens without INH for treating TB disease are less well studied than those with INH. CDC, the American Thoracic Society, and the Infectious Diseases Society of America have recommended a regimen of rifampin, pyrazinamide, and ethambutol for 6 months, with consideration of a fluoroquinolone, such as moxifloxacin or levofloxacin, for more extensive disease, when the infecting *M. tuberculosis* isolate is INH-resistant or the patient is intolerant of INH (4). Directly observed therapy should be used for reducing the potential for acquired rifampin resistance.

An alternative treatment regimen for LTBI, when the infecting *M. tuberculosis* is believed to be INH resistant, or when the patient cannot tolerate INH, is rifampin administered for 4 months, either self-administered or directly observed (5). This regimen has been adopted for routine use by some U.S. TB program directors. A combination regimen of INH and rifampin for 3 or 4 months is used overseas and in some U.S. jurisdictions, although it is not currently recommended in U.S. national guidelines (5,6). This regimen could be prescribed via a fixed dose combination of INH and rifampin if supplies for complete treatment courses are in stock. The combination regimen of rifampin and pyrazinamide for 2 months is *not recommended because of excess toxicity*. Any of the alternative regimens for TB disease or LTBI could be more expensive than the standard regimens.

CDC is working with the Food and Drug Administration (FDA), the National Tuberculosis Controllers Association, and the two pharmaceutical companies currently planning to supply INH (Teva Pharmaceuticals USA, Sellersville, Pennsylvania; and Sandoz Inc., Princeton, New Jersey) to determine current INH stocks, jurisdictions with urgent needs, and options for restoring supplies as soon as can be safely done. CDC will monitor notifications from public health officials of INH shortages and stocks, and will provide updates as they are available on the website for the Division of Tuberculosis Elimination at <http://www.cdc.gov/tb/>.

References for More Information:

1. CDC. Notes from the field: national shortage of isoniazid 300 mg tablets. *MMWR*. 2012;61:1029. Available at http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6150a4.htm?s_cid=mm6150a4_w
2. FDA. Drug safety and availability. Available at <http://www.fda.gov/drugs/drugsafety/drugshortages/default.htm>.
3. World Health Organization. Treatment of tuberculosis: guidelines, fourth edition. WHO Press, 2010. Available at http://whqlibdoc.who.int/publications/2010/9789241547833_eng.pdf.
4. CDC. Treatment of Tuberculosis, American Thoracic Society, CDC, and Infectious Diseases Society of America. *MMWR* 2003;52(No. RR-11). Available at <http://www.cdc.gov/mmwr/PDF/rr/rr5211.pdf>.
5. CDC. Targeted tuberculin testing and treatment of latent tuberculosis infection. *MMWR* 2000;49(No. RR-6), available at <http://www.cdc.gov/mmwr/PDF/rr/rr4906.pdf>.
6. National Institute for Health and Clinical Excellence. Tuberculosis: clinical diagnosis and management of tuberculosis, and measures for its prevention and control. NICE clinical guideline 117. Royal College of Physicians of London, 2006 and 2011. Available at <http://www.nice.org.uk/nicemedia/live/13422/53638/53638.pdf>.

DHEC contact information for reportable diseases and reporting requirements

Reporting of diseases is consistent with South Carolina Law requiring the reporting of diseases and conditions to your state or local public health department. (State Law # 44-29-10 and Regulation # 61-20) as per the DHEC 2013 List of Reportable Conditions available at:

<http://www.scdhec.gov/health/disease/reportables.htm>

Federal HIPAA legislation allows disclosure of protected health information, without consent of the individual, to public health authorities to collect and receive such information for the purpose of preventing or controlling disease. (HIPAA 45 CFR §164.512).

Regional Public Health Offices – 2013

Mail or call reports to the Epidemiology Office in each Public Health Region.

Region 1

Anderson, Oconee

220 McGee Road
Anderson, SC 29625
Phone: (864) 260-4358
Fax: (864) 260-5623
Nights / Weekends: 1-866-298-4442

Abbeville, Edgefield, Greenwood, Laurens, McCormick, Saluda

1736 S. Main Street
Greenwood, SC 29646
Phone: 1-888-218-5475
Fax: (864) 942-3690
Nights / Weekends: 1-800-420-1915

Region 2

Greenville, Pickens

PO Box 2507
200 University Ridge
Greenville, SC 29602-2507
Phone: (864) 282-4139
Fax: (864) 282-4373
Nights / Weekends: (864) 809-3825

Cherokee, Spartanburg, Union

PO Box 4217
151 E. Wood Street
Spartanburg, SC 29305-4217
Phone: (864) 282-4139
Fax: (864) 282-4373
Nights / Weekends: (864) 809-3825

Region 3

Chester, Lancaster, York

PO Box 817
1833 Pageland Highway
Lancaster, SC 29720
Phone: (803) 286-9948
Fax: (803) 286-5418
Nights / Weekends: 1-866-867-3886

Region 3 (continued)

Fairfield, Lexington, Newberry, Richland

2000 Hampton Street
Columbia, SC 29204
Phone: (803) 576-2749
Fax: (803) 576-2993
Nights / Weekends: 1-888-554-9915

Region 4

Chesterfield, Darlington, Dillon, Florence, Marlboro, Marion

145 E. Cheves Street
Florence, SC 29506
Phone: (843) 661-4830
Fax: (843) 661-4859
Nights / Weekends: (843) 601-7051

Clarendon, Kershaw, Lee, Sumter

PO Box 1628
105 North Magnolia Street
Sumter, SC 29150
Phone: (803) 773-5511
Fax: (803) 775-9941
Nights/Weekends: 803-458-1847

Region 5

Bamberg, Calhoun, Orangeburg

PO Box 1126
1550 Carolina Avenue
Orangeburg, SC 29116
Phone: (803) 533-7199
Fax: (803) 533-7134
Nights / Weekends: (803) 535-9265

Aiken, Allendale, Barnwell

222 Beaufort Street, NE
Aiken, SC 29801
Phone: (803) 642-1618
Fax: (803) 643-8386
Nights / Weekends: (803) 645-8167

Region 6

Georgetown, Horry, Williamsburg

1931 Industrial Park Road
Conway, SC 29526-5482
Phone: (843) 915-8804
Fax: (843) 365-0085
Nights/Weekends: (843) 340-4754

Region 7

Berkeley, Charleston, Dorchester

4050 Bridge View Drive, Suite 600
N. Charleston, SC 29405
Phone: (843) 953-0047
Fax: (843) 953-0051
Nights / Weekends: (843) 219-8470

Region 8

Beaufort, Colleton, Hampton, Jasper

219 S. Lemacks Street
Walterboro, SC 29488
Phone: (843) 441-1091
Fax: (843) 549-6845
Nights / Weekends: 1-843-441-1091

DHEC Bureau of Disease Control

Division of Acute Disease Epidemiology

1751 Calhoun Street
Box 101106
Columbia, SC 29211
Phone: (803) 898-0861
Fax: (803) 898-0897
Nights / Weekends: 1-888-847-0902



South Carolina Department of Health
and Environmental Control

www.scdhec.gov

Categories of Health Alert messages:

Health Alert

Conveys the highest level of importance; warrants immediate action or attention.

Health Advisory

Provides important information for a specific incident or situation; may not require immediate action.

Health Update

Provides updated information regarding an incident or situation; unlikely to require immediate action.